

The lepton anomaly in the presence of torsion

Antonio Dobado

Departamento de Física Teórica
Universidad Complutense de Madrid
28040 Madrid, Spain

and

Antonio L. Maroto

Departamento de Física Teórica
Universidad Autónoma de Madrid
28049 Madrid, Spain

February 1, 2008

Abstract

In the presence of a curved space-time with torsion, the divergence of the leptonic current gets new contributions from the curvature and the torsion in addition to those coming from the electroweak fields. However it is possible to define a new lepton current which is gauge and local Lorentz invariant whose divergence gets no contribution from the torsion. Therefore we argue that torsion does not contribute to anomalous lepton number production.

One of the most important open problems in modern physics is that of the origin of the baryon and lepton asymmetries of the universe. From the standard big-bang nucleosynthesis and the present data it is possible to set the value of the baryon versus photon densities in something about $n_B/n_\gamma \simeq 10^{-10}$. Since today the antibaryon density is thought to be completely negligible ($n_B \gg n_{\bar{B}}$), the above ratio can be considered also as the value of the baryon asymmetry (δ) before the decoupling of baryons and antibaryons (in fact quarks and antiquarks) and radiation, i.e. $\delta = (n_B - n_{\bar{B}})/(n_B + n_{\bar{B}}) \simeq 10^{-10}$. The main problem of any baryogenesis model is to explain the origin of this small number which however probably gives rise to most of the observed matter around us today.

Sometime ago Sakharov [1] set some necessary ingredients in order to get baryon asymmetry starting from a symmetric universe at early times. This asymmetry production requires obviously B (baryon number) violation but also C and CP violation and absence of thermal equilibrium. The first modern models of baryogenesis appeared last decade with the advent of Grand Unified Theories (GUT) (see for example [2] for a review). In these models the decays of the X and Y gauge or the Higgs bosons are B violating. On the other hand C and CP breaking is provided by chiral interactions and complex Higgs selfcouplings in a natural way. The absence of thermal equilibrium occurs whenever the decay rate of the above mentioned bosons is smaller than the expansion rate of the universe.

More recently it was pointed out that the anomalous electroweak L (lepton) and B numbers non-conservation present in the Standard Model (SM) [3], which preserves $B - L$, could be enough to wash out any B or L asymmetry generated at the GUT scale unless the universe started with some non-vanishing $B - L$ [4]. In that work the authors studied also the possibility of baryon asymmetry production by these anomalous electroweak processes themselves but they arrived to the conclusion that this is not possible if the electroweak phase transition is of second order as it seems to be the case.

When the gravitational field is taken into account in the computation of the L and B

number anomalies [5], one observes that $B - L$ is not conserved any more. Thus one could speculate on the possibility that these gravitational effects could generate a $B - L$ asymmetry and that the usual electroweak effect would just redistribute the relative amounts of B and L . However, it is not obvious how to make such an scenario to work in detail [6].

In this work we will consider another potential source of anomalous $B - L$ non-conservation which could appear when, in addition to the gravitational field, we have also torsion. As it is well known, once a metric is given, there is a unique affine connection which is metric compatible and torsion free. This is the Levi-Civita connection which is given by the Christoffel symbols and it was the only one considered by Einstein in his original formulation of General Relativity. However, in all the modern discussions on gravitation, the metric (the vierbein) and the connection are treated as independent structures and then torsion appears in a natural way. In particular, whenever fermions are present, as it happens in the SM, the torsion is different from zero [7]. In addition, when one considers only minimal couplings, fermions are the only SM particles which couple to the torsion, in particular, to its pseudo-trace. In a recent work [8] the authors have computed the B and L anomaly for the SM defined on a gravitational field with torsion by which we mean a metric (vierbein) and an arbitrary (but metric compatible) affine connection. We have found that, in addition to the electroweak and curvature terms, there are also new contributions coming from torsion to the L anomaly. Thus, any amount of torsion present in the background gravitational field can in principle be transformed in lepton number. This mechanism for the $B - L$ production could be used to build a new scenario of lepton asymmetry generation. The resulting lepton asymmetry could be transformed again by the more usual electroweak anomaly effects in a baryon asymmetry. In the following we will discuss that issue in detail and we will conclude that, due to the possibility of redefining a gauge and local Lorentz invariant lepton current in presence of torsion, no real anomalous lepton production takes place from the above mechanism.

The Euclidean SM matter lagrangian in a curved space-time with torsion can be written

for one family as follows [8]:

$$\mathcal{L}_m = Q^\dagger \not{D}^Q Q + L^\dagger \not{D}^L L \quad (1)$$

where the matter fields appear in doublets:

$$Q = \begin{bmatrix} u \\ d \end{bmatrix} \quad L = \begin{bmatrix} \nu \\ e \end{bmatrix} \quad (2)$$

The Dirac operators for quarks and leptons are given by

$$\begin{aligned} i\not{D}^Q &= i\gamma^\mu(\partial_\mu + \boldsymbol{\Omega}_\mu^Q + \mathbf{G}_\mu + \mathbf{W}_\mu^Q P_L + \mathbf{B}_\mu^Q + \mathbf{S}_\mu^Q \gamma_5) \\ i\not{D}^L &= i\gamma^\mu(\partial_\mu + \boldsymbol{\Omega}_\mu^L + \mathbf{W}_\mu^L P_L + \mathbf{B}_\mu^L + \mathbf{S}_\mu^L \gamma_5) \end{aligned} \quad (3)$$

Here $\mathbf{G}_\mu = -ig_S G_\mu^a \Lambda^a$, $\mathbf{W}_\mu = -ig W_\mu^a T^a$ and $\mathbf{B}_\mu = ig' B_\mu (\mathbf{y}_L P_L + \mathbf{y}_R P_R)$ are the usual gauge fields corresponding to the $SU(3)_c$, $SU(2)_L$ and $U(1)_Y$ groups respectively and the hypercharge matrices are given by

$$\mathbf{y}_{L,R}^{Q,L} = \begin{pmatrix} y_{L,R}^{u,\nu} & \\ & y_{L,R}^{d,e} \end{pmatrix} \quad (4)$$

In addition we have the Levi-Civita spin connection and torsion terms that for leptons read:

$$\boldsymbol{\Omega}_\mu^L = -\frac{i}{2} \Gamma_\mu^{ab} \begin{pmatrix} P_L \Sigma_{ab} & \\ & \Sigma_{ab} \end{pmatrix}, \quad \mathbf{S}_\mu^L \gamma_5 = -\frac{1}{8} S_\mu \begin{pmatrix} P_L \gamma_5 & \\ & \gamma_5 \end{pmatrix} \quad (5)$$

where Σ_{ab} are the $SO(4)$ generators in the spinor representation, $\{\Gamma_\mu^{ab}\}$ are the components of the Levi-Civita spin connection and $S_\alpha = \epsilon_{\mu\nu\lambda\alpha} T^{\mu\nu\lambda}$ is the pseudo-trace of the torsion tensor. Such a tensor is defined as follows: given a metric connection $\{\hat{\Gamma}_\mu^{ab}\}$, it can be written in general as:

$$\hat{\Gamma}_\mu^{ab} = \Gamma_\mu^{ab} + e_\nu^a e^{\lambda b} K_{\mu\lambda}^\nu \quad (6)$$

where e_ν^a is the vierbein and $K_{\mu\lambda}^\nu$ is known as the contorsion tensor which is related to the torsion tensor by [9]:

$$K^{\nu\mu\lambda} = \frac{1}{2} (T^{\nu\mu\lambda} + T^{\mu\nu\lambda} + T^{\lambda\nu\mu}) \quad (7)$$

Since there are no right neutrinos, the upper components in Eq.5 appear multiplied by a left projector. In the case of quarks such projectors are not present. As is well known, the above lagrangian is invariant under the following global transformations:

$$Q \rightarrow Q + i\alpha Q, L \rightarrow L + i\beta L \quad (8)$$

The corresponding Noether currents are nothing but the baryon and lepton number currents:

$$j_B^\mu = \frac{1}{N_c} Q^\dagger \gamma^\mu Q, j_L^\mu = L^\dagger \gamma^\mu L \quad (9)$$

which classically are conserved, i.e $\nabla_\mu j_{Q,L}^\mu = 0$, where ∇_μ is the Levi-Civita covariant derivative. However these conservation laws are violated due to quantum effects. The corresponding anomalies can be obtained using, for instance, the Fujikawa method based on the $i\cancel{D}^\dagger i\cancel{D}$ and $i\cancel{D} i\cancel{D}^\dagger$ operators [10]. The gaussian regulators associated to these operators respect the gauge and local Lorentz symmetries of the theory. This procedure yields for the anomalies the following results (for Euclidean signature) [8]:

$$\nabla_\mu j_B^\mu = \frac{1}{32\pi^2} \epsilon^{\mu\nu\alpha\beta} \left(\frac{g^2}{2} W_{\mu\nu}^a W_{\alpha\beta}^a + g'^2 B_{\mu\nu} B_{\alpha\beta} \sum_{u,d} (y_L^2 - y_R^2) \right) \quad (10)$$

and

$$\begin{aligned} \nabla_\mu j_L^\mu = \frac{1}{32\pi^2} \left\{ -\frac{\epsilon^{\alpha\beta\gamma\delta}}{24} R_{\mu\nu\alpha\beta} R^{\mu\nu}_{\gamma\delta} + \frac{\epsilon^{\alpha\beta\gamma\delta}}{48} S_{\beta;\gamma} S_{\delta;\alpha} + \epsilon^{\alpha\beta\gamma\delta} \left(\frac{g^2}{2} W_{\gamma\delta}^a W_{\alpha\beta}^a \right. \right. \\ \left. \left. + g'^2 B_{\gamma\delta} B_{\alpha\beta} \sum_{\nu,e} (y_L^2 - y_R^2) \right) + \frac{1}{6} \square S_{;\alpha}^\alpha + \frac{1}{96} (S^\alpha S^\nu S_\alpha)_{;\nu} - \frac{1}{6} \left(R^{\nu\alpha} S_\alpha - \frac{1}{2} R S^\nu \right)_{;\nu} \right\} \quad (11) \end{aligned}$$

These expressions are gauge and local Lorentz invariant. In principle, it is possible to use a different regulator within the Fujikawa method and, in fact, the $(i\cancel{D})^2$ operator is also frequently used in the calculation of the so called consistent gauge anomaly. This regulator yields no anomaly in L and B but its eigenvalues are not gauge invariant. The choice of regulator have the same effects as the choice of the shifted momenta in the divergent integrals appearing in the triangle diagrams contributing to the axial anomaly. In that case, it is

possible to choose the integrated momenta in such a way that the axial anomaly vanishes, but this introduces an anomaly in the gauge current. Conversely, it is possible to cancel the gauge anomaly but maintaining the axial one. However axial anomalies have measurable physical effects as the π^0 decay into two photons and therefore the above ambiguities in their precise value should be removed. This can be achieved by demanding that gauge invariance is respected in the calculations which in turn imposes the use of a gauge and local Lorentz invariant regulator as the one considered here to obtain Eq.10 and Eq.11.

The resulting lepton anomaly has terms depending on the curvature and the torsion that appear due to the absence of one of the chirality components of the neutrino field. Such terms are not present in the baryonic anomaly since quarks have both chirality components. Thus in contrast with flat space-time, $B - L$ is spoiled in the presence of curvature and in principle also in presence of torsion.

Therefore we have three different kinds of potential contributions to the lepton anomaly. First we have a $\epsilon^{\alpha\beta\gamma\delta} R_{\mu\nu\alpha\beta} R^{\mu\nu}_{\gamma\delta}$ term where $R^{\mu\nu}_{\gamma\delta}$ is the curvature associated to the Levi-Civita spin connection. This term integrated over the whole space-time is related to a topological invariant called the Pontryagin index of the manifold [11] and it can be written as a four-divergence, i.e, $\nabla_\mu K^\mu = \epsilon^{\alpha\beta\gamma\delta} / (768\pi^2) R_{\mu\nu\alpha\beta} R^{\mu\nu}_{\gamma\delta}$. This fact allows us to redefine the lepton current as $\tilde{j}_L^\mu = j_L^\mu + K^\mu$. In absence of the other two pieces of the anomaly, this current would be conserved, although it is not local Lorentz invariant. Such a topological term could give rise to actual contributions to the L violation through the so called gravitational instantons [12] that would be relevant in the context of quantum gravity and, as mentioned above, its possible relevance for the generation of asymmetries was sketched in [6]. Second, there is also the well known $SU(2)_L$ and $U(1)_Y$ gauge contributions to the anomaly which were studied in a cosmological context in [4]. Finally, we also have the possible torsion contributions in which we are interested in this work.

Concerning to the possibility of having anomalous lepton production from torsion two

important facts should be mentioned. First one could think, following the usual topological interpretation of the axial anomalies, that the lepton number anomaly could be given by the index of the Dirac operator with a spin connection with torsion. In other words: $\nabla_\mu j_L^\mu = (\epsilon^{\alpha\beta\gamma\delta}/(768\pi^2)\hat{R}_{\mu\nu\alpha\beta}\hat{R}^{\mu\nu}_{\gamma\delta} + \{\text{electroweak gauge terms}\})$ where $\hat{R}_{\mu\nu\alpha\beta}$ is the curvature associated to the full connection (with torsion). However it can be explicitly shown that this is not the case since for a completely antisymmetric torsion we have

$$\begin{aligned} \frac{\epsilon^{\alpha\beta\gamma\delta}}{24}\hat{R}_{\mu\nu\alpha\beta}\hat{R}^{\mu\nu}_{\gamma\delta} &= \frac{\epsilon^{\alpha\beta\gamma\delta}}{24}R_{\mu\nu\alpha\beta}R^{\mu\nu}_{\gamma\delta} + \frac{1}{432}\epsilon^{\alpha\beta\gamma\delta}S_{\alpha;\beta}S_{\gamma;\delta} \\ &+ \frac{1}{6}\left(\frac{1}{3}R^{\beta\alpha}S_{\alpha;\beta} - \frac{1}{6}S^\mu_{;\mu}R - \frac{1}{432}S^2S^\mu_{;\mu} - \frac{1}{216}S_{\alpha;\beta}S^\alpha S^\beta\right) \end{aligned} \quad (12)$$

and this expression does not agree with the result in Eq.11.

Second, and more relevant is the fact that the torsion contribution to the anomaly is a four-divergence and it can be absorbed in the redefinition of the lepton current as follows:

$$\tilde{j}_L^\mu = j_L^\mu - \frac{1}{32\pi^2}\left(\frac{1}{6}S^\alpha_{;\alpha\mu} + \frac{1}{96}S^\alpha S^\mu S_\alpha - \frac{1}{6}\left(R^{\mu\alpha}S_\alpha - \frac{1}{2}RS^\mu\right) + \frac{1}{48}\epsilon^{\mu\beta\gamma\delta}S_{\beta;\gamma}S_\delta\right) \quad (13)$$

The new current $\tilde{j}_L^\mu$ is still gauge and local Lorentz invariant and depends explicitly on the torsion. In the absence of the other contributions to the anomaly this current is conserved, that is, $\nabla_\mu \tilde{j}_L^\mu = 0$. Thus we observe that the possible effects of the torsion can be eliminated away by this redefinition of the lepton current. Notice that the new definition, being gauge and local Lorentz invariant, is physically meaningful and should satisfy the appropriate Ward identities for a properly defined lepton current.

Let us stress once again the main differences between the torsion terms and the gauge and gravitational contributions to the lepton anomaly. First, as we have just mentioned, the redefined lepton current in Eq.13 is gauge and local Lorentz invariant, moreover there is no local symmetry associated to torsion that is spoiled in the redefinition. However when one absorbs in a similar way the gauge and gravitational terms, the new lepton current no longer respects gauge and Lorentz symmetries. In addition, the gauge and gravitational

terms appear in the form of topological invariants. This fact implies that the variation of the lepton number is related to quantum tunneling processes between different classical vacua characterized by different winding numbers. Such processes, as it is well known, are dominated by instantons in the semi-classical approximation. However, as we mentioned before, the torsion contribution is not topological. In fact it has been shown [13] that apart from the Pontryagin and Euler classes, there is only one more non-trivial 4-form containing torsion and such form does not agree with the result in Eq.11.

An alternative way to show that torsion does not contribute to the lepton anomaly is based on a different choice of regulator. In fact let us take as regulators for the anomalies the operators $i\bar{\mathcal{D}}^\dagger i\bar{\mathcal{D}}$ and $i\bar{\mathcal{D}}i\bar{\mathcal{D}}^\dagger$, where $\bar{\mathcal{D}}$ is the $\mathcal{D}$ operator in which the torsion field has been set to zero. These operators respect all the gauge and local Lorentz symmetries of the theory and therefore they are also valid as regulators. However they do not depend on torsion, this implies that the regulated anomalies cannot depend on torsion, as expected.

To summarize, it is shown that, although there are torsion contributions to the lepton anomaly (which has been obtained by means of a gauge and local Lorentz invariant regulator), however such terms cannot give rise to lepton number generation.

Acknowledgments: This work has been supported in part by the Ministerio de Educación y Ciencia (Spain) (CICYT AEN96-1634).

References

- [1] A.D. Sakharov, *Zh. Eksp. Teor. Fiz. Pis'ma* **5** (1967) 32 (transl. *JETP Lett.* **5** (1967) 24)
- [2] E. W. Kolb and M.S. Turner *Ann. Rev. Nucl. Part. Sci.* **33** (1983) 645
- [3] G. 't Hooft, *Phys. Rev. Lett.* **37** (1976) 8; *Phys. Rev.* **D14** (1976) 3432
- [4] V.A. Kuzmin, V.A. Rubakov and M.E. Shaposhnikov, *Phys. Lett.* **155B** (1985) 36
- [5] R. Delbourgo and A. Salam, *Phys. Lett.* **B40** (1972) 381
- [6] L. Ibáñez, *Proceedings of the International Europhysics Conference on High Energy Physics*, Marseille (1993). Editions Frontieres.
- [7] F. W. Hehl, P. von der Heyde, G. D. Kerlick and J. M. Nester, *Rev. Mod. Phys.* **48**, (1976), 393
- [8] A. Dobado and A.L. Maroto, *Phys. Rev.* **D54** (1996) 5185
- [9] M. Nakahara, *Geometry, Topology and Physics*, IOP Publishing, 1990
- [10] K. Fujikawa, *Phys. Rev.* **D21** (1980) 2848; *ibid* **D29** (1984) 285
- [11] T. Eguchi, P.B. Gilkey and A.J. Hanson, *Phys. Rep.* **66** (1980) 213
- [12] T. Eguchi and P.G.O. Freund, *Phys. Rev. Lett.* **37** (1976) 1251
- [13] H. T. Nieh and M.L. Yan, *J. Math. Phys.* **23** (1982) 373